## A Thermal Map for all Oregon Streams

















Funded by:

The aquatics community within Oregon has amassed significant amounts of stream temperature data through their collective monitoring efforts in previous decades. As part of a broader regional effort, the NorWeST project has developed a comprehensive interagency stream temperature database for Oregon that consists of 18,774 summers of monitoring effort at 6,461 unique sites (map inset). Those data were used with spatial statistical network models (details at the SSN/STARS website:

www.fs.fed.us/rm/boise/AWAE/projects/

SpatialStreamNetworks.shtml) to develop an accurate stream temperature model (R² = 90%; RMSE = 1.0°C), which was then used to predict high resolution (1 kilometer) historical and future climate scenarios for all of Oregon's streams. This poster depicts a historical scenario of the mean August temperature from 1993-2011 mapped to the USGS 1:100,000-scale NHDPlus hydrography layer trimmed to exclude reaches >15% slope. NorWeST stream climate scenarios are available as ArcGIS shapefiles from the project website (www.fs.fed.us/rm/boise/AWAE/projects/ NorWeST.html) and can also be viewed dynamically online (www.sciencebase.gov/flexviewer/NorWeST/). Daily summaries (min/max/mean) of the temperature data used to develop the temperature model are available through the website if permission was given for their distribution. All data are attributed to the original source agency and contributing biologists/hydrologists in metadata files.

By providing open access to stream temperature information in user-friendly formats, the NorWeST project is facilitating coordination of monitoring activities among organizations, better conservation planning, and new research on temperature dynamics and thermal ecology. The analytical infrastructure used to develop the Oregon stream temperature model consists of a new class of spatial statistical model for data on stream networks that could also be used with water chemistry attributes (e.g., pH, alkalinity, conductivity, etc.), various biological datasets (species occurrence, abundance, genetic attributes), or habitat surveys to provide a wealth of new information about streams.

## KEY REFERENCES

NorWeST Website: www.fs.fed.us/rm/boise/AWAE/projects/ NorWeST.html

## **Spatial Statistical Network Models**

Isaak, D., C. Luce, B. Rieman, D. Nagel, E. Peterson, D. Horan, S. Parkes, G. Chandler. 2010. Effects of climate change and wildfire on stream temperatures and salmonid thermal habitat in a mountain river network. Ecological Applications 20: 1350-1371.

Isaak, D., E. Peterson, J. Ver Hoef, S. Wenger, J. Falke, C. Torgersen, C. Sowder, A. Steel, M. Fortin, C. Jordan, A. Reusch, N. Som, P. Monestiez. 2014. Applications of spatial statistical network models to stream data. Wiley Interdisciplinary Reviews -Water 1:277-294.

Peterson, E., J. Ver Hoef, D. Isaak, J. Falke, M. Fortin, C. Jordan, K. McNyset, P. Monestiez, A. Ruesch, A. Sengupta, N. Som, E. Steel, D. Theobald, C. Torgersen, and S. Wenger. 2013. Modeling dendritic ecological networks in space: an integrated network perspective. Ecology Letters 16:707-719.

Ver Hoef, J., E. Peterson, and D. Theobald. 2006. Spatial statistical models that use flow and stream distance. Environmental and Ecological Statistics 13:449-464.

Ver Hoef, J., E. Peterson, D. Clifford, and R. Shah. 2014. SSN: An R package for spatial statistical modeling on stream networks. Journal of Statistical Software **56(3)**:1-47.

